

## Sinus Lift or Augmentation Surgery | Post Operative Instructions

These are additional instructions following a Sinus Lift or Augmentation Surgery. Please review our General Post-Operative instructions in addition to the following information that applies when the upper jawbone height or width have been lost. The graft is placed to help restore your jawbone in preparation for possible implant replacement of the missing tooth or teeth.

You have had a Sinus Lift Augmentation procedure in your upper jaw. This procedure regains lost bone height in the area of your first and second molar and occasionally second premolar. It is an important procedure as it allows implant placement in an area that could not be implanted otherwise because of insufficient bone height due to an enlarged sinus.

The bone that has been grafted is most commonly a combination of freeze-dried bone, artificial synthetic bone and your own bone. Because of this you may have two post-surgical wounds, the donor site and the recipient site.

**DO NOT UNDER ANY CIRCUMSTANCES BLOW YOUR NOSE FOR THE NEXT FOUR (4) WEEKS.** This may be longer if indicated. You may sniff all you like but **NO BLOWING**.

Do not blow your nose or sneeze holding your nose. Sneeze with your mouth open. Do not drink with straws and do not spit. Scuba diving and flying in pressurized aircraft may also increase sinus pressure and should be avoided. Decongestants such as *Drixoral*, *Dimetapp* or *Sudafed* will reduce pressure in the sinuses. You may also be given a prescription for antibiotics. Please take them as directed. Anything that causes pressure in the nasal cavity must be avoided. Avoid “bearing down”—as when lifting heavy objects, blowing up balloons, playing musical instruments that require a blowing action or any other activity that increases nasal or oral pressure. Smoking must be stopped.

### **Antibiotics**

Be sure to take the prescribed antibiotic as directed to help prevent infection.

### **Oral Hygiene**

Do not rinse on the day of your surgery. This tends to disturb the blood clot, and open the wound and can prolong bleeding and slow the healing process. You should not have a significant amount of blood in your mouth. Saliva can be swallowed, even if slightly blood tinged.

Keeping your mouth clean after surgery is essential to reduce the risk of infection. Start salt water rinses the day following your procedure, Use one-half teaspoon of salt dissolved in an 8 ounce glass of warm water and gently rinse with portions of the solution, taking five minutes to use the entire glassful. Repeat as often as you like, but at least four to five times daily and always after eating for the next five days.

Do not brush the teeth in the area of the surgery for 48 hours. When brushing, be very gentle. Also, be very gentle when you are expectorating.

We may prescribe an antibiotic rinse (*Chlorhexadine, Periogard, and Peridex*) for certain procedures. This rinse should be used in the morning and at bedtime after routine mouth care. Do not eat or rinse your mouth after using the medicated rinse. Using this rinse more than two times a day will cause staining of your teeth.

### **Smoking**

Do not smoke for at least two weeks after surgery, if at all. As discussed at your consultation smoking dramatically increases the risk of gone graft and sinus augmentation failure.

### **Wearing your Prosthesis or Night Guards**

Partial dentures, flippers, or full dentures should not be used immediately after surgery until your post-operative appointment unless specifically instructed otherwise. Please contact the office if there is any question. If you have a temporary “flipper” to wear do not place it until the numbness in the area is gone. When it is placed it should not touch the gums in the area of the surgery. If it does, this can cause ulceration of the wound edges and breakdown the suture margins. This can lead to loss of the graft. If you have any questions about the fit of your flipper, partial or complete denture, do not wear it until Dr. Deck can see you.

### **Post-Operative Problems or Complications**

As with any procedure, unexpected post-operative healing can occur, if you notice the unexpected flow of air or liquids between your mouth and nose, please let us know immediately.

- If you are aware of several small partials of graft material being discharged from your nose, let us know!
- If you experience sinus or nasal congestion on the side your surgery was performed, let us know!
- If there is an increase in swelling in your mouth in your mouth, cheek or under your eye after 3 days, let us know!

It is our desire that your recovery be as smooth and pleasant as possible. Following these instructions will assist you, but if you have any questions about you progress, please call the office at 519 348 8808